

Set Operations

Section 2.2



Section Summary

- Set Operations
 - Union
 - Intersection
 - Complementation
 - Difference
- More on Set Cardinality
- Set Identities
- Proving Identities
- Membership Tables



Boolean Algebra

- Propositional calculus and set theory are both instances of an algebraic system called a *Boolean Algebra*. This is discussed in Chapter 12.
- The operators in set theory are analogous to the corresponding operator in propositional calculus.
- As always there must be a universal set U . All sets are assumed to be subsets of U .

Union


- **Definition:** Let A and B be sets. The *union* of the sets A and B , denoted by $A \cup B$, is the set:

$$\{x \mid x \in A \vee x \in B\}$$

- **Example:** What is $\{1,2,3\} \cup \{3,4,5\}$?

Solution: $\{1,2,3,4,5\}$

Venn Diagram for $A \cup B$



Intersection

- **Definition:** The *intersection* of sets A and B , denoted by $A \cap B$, is

$$\{x \mid x \in A \wedge x \in B\}$$


- Note if the intersection is empty, then A and B are said to be *disjoint*.
- **Example:** What is? $\{1,2,3\} \cap \{3,4,5\}$?

Solution: $\{3\}$

- **Example:** What is?
 $\{1,2,3\} \cap \{4,5,6\}$?

Solution: \emptyset

Venn Diagram for $A \cap B$



Complement

Definition: If A is a set, then the complement of the A (with respect to U), denoted by \bar{A} is the set $U - A$


$$\bar{A} = \{x \in U \mid x \notin A\}$$

(The complement of A is sometimes denoted by A^c .)

Example: If U is the positive integers less than 100, what is the complement of $\{x \mid x > 70\}$

Solution: $\{x \mid x \leq 70\}$


Venn Diagram for Complement



Difference

- **Definition:** Let A and B be sets. The *difference* of A and B , denoted by $A - B$, is the set containing the elements of A that are not in B . The difference of A and B is also called the complement of B with respect to A .

$$A - B = \{x \mid x \in A \wedge x \notin B\} = A \cap \bar{B}$$




Venn Diagram for $A - B$

The Cardinality of the Union of Two Sets

- Inclusion-Exclusion

$$|A \cup B| = |A| + |B| - |A \cap B|$$



Venn Diagram for $A, B, A \cap B, A \cup B$

- **Example:** Let A be the math majors in your class and B be the CS majors. To count the number of students who are either math majors or CS majors, add the number of math majors and the number of CS majors, and subtract the number of joint CS/math majors.
- We will return to this principle in Chapter 6 and Chapter 8 where we will derive a formula for the cardinality of the union of n sets, where n is a positive integer.

Review Questions

Example: $U = \{0,1,2,3,4,5,6,7,8,9,10\}$ $A = \{1,2,3,4,5\}$, $B = \{4,5,6,7,8\}$

1. $A \cup B$

Solution: $\{1,2,3,4,5,6,7,8\}$

2. $A \cap B$

Solution: $\{4,5\}$

3. \bar{A}

Solution: $\{0,6,7,8,9,10\}$

4. \bar{B}

Solution: $\{0,1,2,3,9,10\}$

5. $A - B$

Solution: $\{1,2,3\}$

6. $B - A$

Solution: $\{6,7,8\}$

Symmetric Difference (*optional*)

Definition: The *symmetric difference* of **A** and **B**, denoted by $A \oplus B$ is the set

$$(A - B) \cup (B - A)$$


Example:

$$U = \{0,1,2,3,4,5,6,7,8,9,10\}$$

$$A = \{1,2,3,4,5\} \quad B = \{4,5,6,7,8\}$$

What is $A \oplus B$:

- **Solution:** $\{1,2,3,6,7,8\}$



Venn Diagram

Set Identities

- Identity laws

$$A \cup \emptyset = A \quad A \cap U = A$$

- Domination laws

$$A \cup U = U \quad A \cap \emptyset = \emptyset$$

- Idempotent laws

$$A \cup A = A \quad A \cap A = A$$

- Complementation law

$$\overline{(\overline{A})} = A$$

Continued on next slide →

Set Identities

- Commutative laws

$$A \cup B = B \cup A \quad A \cap B = B \cap A$$

- Associative laws

$$A \cup (B \cup C) = (A \cup B) \cup C$$

$$A \cap (B \cap C) = (A \cap B) \cap C$$

- Distributive laws

$$A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$$

$$A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$$

Continued on next slide →

Set Identities

- De Morgan's laws

$$\overline{A \cup B} = \overline{A} \cap \overline{B} \quad \overline{A \cap B} = \overline{A} \cup \overline{B}$$

- Absorption laws

$$A \cup (A \cap B) = A \quad A \cap (A \cup B) = A$$

- Complement laws

$$A \cup \overline{A} = U \quad A \cap \overline{A} = \emptyset$$

Proving Set Identities

- Different ways to prove set identities:
 1. Prove that each set (side of the identity) is a subset of the other.
 2. Use set builder notation and propositional logic.
 3. Membership Tables: Verify that elements in the same combination of sets always either belong or do not belong to the same side of the identity. Use 1 to indicate it is in the set and a 0 to indicate that it is not.

Proof of Second De Morgan Law

Example: Prove that $\overline{A \cap B} = \overline{A} \cup \overline{B}$

Solution: We prove this identity by showing that:

$$1) \quad \overline{A \cap B} \subseteq \overline{A} \cup \overline{B} \quad \text{and}$$

$$2) \quad \overline{A} \cup \overline{B} \subseteq \overline{A \cap B}$$

Continued on next slide →

Proof of Second De Morgan Law

These steps show that: $\overline{A \cap B} \subseteq \overline{A} \cup \overline{B}$

$$x \in \overline{A \cap B}$$

$$x \notin A \cap B$$

$$\neg((x \in A) \wedge (x \in B))$$

$$\neg(x \in A) \vee \neg(x \in B)$$

$$x \notin A \vee x \notin B$$

$$x \in \overline{A} \vee x \in \overline{B}$$

$$x \in \overline{A} \cup \overline{B}$$

by assumption

defn. of complement

defn. of intersection

1st De Morgan Law for Prop Logic

defn. of negation

defn. of complement

defn. of union

Continued on next slide →

Proof of Second De Morgan Law

These steps show that:

$$\overline{A} \cup \overline{B} \subseteq \overline{A \cap B}$$

$$x \in \overline{A} \cup \overline{B}$$

$$(x \in \overline{A}) \vee (x \in \overline{B})$$

$$(x \notin A) \vee (x \notin B)$$

$$\neg(x \in A) \vee \neg(x \in B)$$

$$\neg((x \in A) \wedge (x \in B))$$

$$\neg(x \in A \cap B)$$

$$x \in \overline{A \cap B}$$

by assumption

defn. of union

defn. of complement

defn. of negation

by 1st De Morgan Law for Prop Logic

defn. of intersection

defn. of complement



Set-Builder Notation: Second De Morgan Law

$$\begin{aligned}\overline{A \cap B} &= \{x \mid x \notin A \cap B\} && \text{by defn. of complement} \\ &= \{x \mid \neg(x \in (A \cap B))\} && \text{by defn. of does not belong symbol} \\ &= \{x \mid \neg(x \in A \wedge x \in B)\} && \text{by defn. of intersection} \\ &= \{x \mid \neg(x \in A) \vee \neg(x \in B)\} && \text{by 1st De Morgan law} \\ &&& \text{for Prop Logic} \\ &= \{x \mid x \notin A \vee x \notin B\} && \text{by defn. of not belong symbol} \\ &= \{x \mid x \in \overline{A} \vee x \in \overline{B}\} && \text{by defn. of complement} \\ &= \{x \mid x \in \overline{A} \cup \overline{B}\} && \text{by defn. of union} \\ &= \overline{A} \cup \overline{B} && \text{by meaning of notation}\end{aligned}$$



Membership Table

Example: Construct a membership table to show that the distributive law holds.

$$A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$$

Solution:

A	B	C	$B \cap C$	$A \cup (B \cap C)$	$A \cup B$	$A \cup C$	$(A \cup B) \cap (A \cup C)$
1	1	1	1	1	1	1	1
1	1	0	0	1	1	1	1
1	0	1	0	1	1	1	1
1	0	0	0	1	1	1	1
0	1	1	1	1	1	1	1
0	1	0	0	0	1	0	0
0	0	1	0	0	0	1	0
0	0	0	0	0	0	0	0

Generalized Unions and Intersections

- Let A_1, A_2, \dots, A_n be an indexed collection of sets.

We define:

$$\bigcup_{i=1}^n A_i = A_1 \cup A_2 \cup \dots \cup A_n$$

$$\bigcap_{i=1}^n A_i = A_1 \cap A_2 \cap \dots \cap A_n$$

These are well defined, since union and intersection are associative.

- For $i = 1, 2, \dots$, let $A_i = \{1, 2, 3, \dots, i\}$. Then,

$$\bigcup_{i=1}^{\infty} A_i = \bigcup_{i=1}^{\infty} \{1, 2, 3, \dots, i\} = \{1, 2, 3, \dots\} = \mathbf{Z}^+$$

$$\bigcap_{i=1}^{\infty} A_i = \bigcap_{i=1}^{\infty} \{1, 2, 3, \dots, i\} = \{1\}.$$